





#### **OPEN QUESTIONS**




#### **OPEN QUESTIONS**































#### Multimessenger Analysis: The Ingredients






# GW and neutrino detectors are "all-sky"




"all-sky" GW+v search

# Neutrinos allow better pointing for EM follow-up searches



prompt search for significant events






## SENSITIVITY TIMELINE

Aasi et al. 2013


	Estimated	$E_{\rm GW} = 10^{-2} M_{\odot} c^2$				Number	% BNS Localized	
	Run	Burst Range (Mpc)		BNS Range (Mpc)		of BNS	within	
Epoch	Duration	LIGO	Virgo	LIGO	Virgo	Detections	$5  \mathrm{deg}^2$	$20  \mathrm{deg}^2$
2015	3 months	40 - 60	_	40 - 80	_	0.0004 - 3	_	_
2016-17	6 months	60 - 75	20 - 40	80 - 120	20 - 60	0.006 - 20	2	5 - 12
2017–18	9 months	75 - 90	40 - 50	120 - 170	60 - 85	0.04 - 100	1 - 2	10 - 12
2019+	(per year)	105	40 - 80	200	65 - 130	0.2 - 200	3 - 8	8 - 28
2022+ (India)	(per year)	105	80	200	130	0.4 - 400	17	48

### MULTIMESSENGER ANALYSIS: THE INGREDIENTS




# Cataloging on the fly

- Can we make a catalog in the right time frame, distance range and sky area?
  - ✓ 1 week
  - ✓ 200-500 Mpc
  - √ 100 deg²
- Extended H-alpha survey (R-band comparison)
  - We only want to find galaxies within horizon distance
  - We don't necessarily need more info than this as long as catalog is complete
- Meter class telescopes work.
- Don't need very high completeness (Hanna+ 2014)




## astrophysical search optimization




Coincidence time window: 500s

# GRB EMISSION EPISODES (PRECURSORS)




Automated survey of BATSE, Fermi and Swift GRBs (2710)
Conclusion --- precursors likely from same central engine activity



## **N**EUTRINO — GAMMA-RAY CORRELATION

 $10^{-3}$ 




Gamma-ray --- neutrino emission mechanism is connected → temporal correlation

--- GRB fluence & neutrino fluence linearly correlated (117 GRBs, from Hummer+ PRL 2012)

Using temporal correlation can decrease False Alarm Rate by x100

Discovery potential for GRBs = 1 TeV neutrino ~ few GeV neutrino




 $10^{-2}$ 


FAR / FAR


 $10^{-1}$ 


Bartos+ PRL 2013

Bartos, Marka PRD(R) 2014












### DETECTION PROSPECTS: ICECUBE+DEEPCORE



### Expanding the GW horizon distance with astrophysics



### Beyond the GW horizon distance with astrophysics



1.5


Neutron star mass  $[M_{\odot}]$ 

2.0

2.5

1.0

0.5



Detection rate improvement
All-sky 14%
External trigger 61%

