

ORIGINAL ARTICLE

IS5 inserts upstream of the master motility operon *flhDC* in a quasi-Lamarckian way

Xiaoxue Wang and Thomas K Wood

Department of Chemical Engineering, Texas A & M University, College Station, TX, USA

Mutation rates may be influenced by the environment. Here, we demonstrate that insertion sequence IS5 in *Escherichia coli* inserts into the upstream region of the *flhDC* operon in a manner that depends on whether the environment permits motility; this operon encodes the master regulator of cell motility, FlhDC, and the IS5 insertion increases motility. IS5 inserts upstream of *flhD*⁺ when cells are grown on soft-agar plates that permit swimming motility, but does not insert upstream of this locus on hard-agar plates that do not permit swimming motility or in planktonic cultures. Furthermore, there was only one IS5 insertion event on soft-agar plates, indicating insertion of IS5 into *flhDC* is not due to general elevated IS5 transposition throughout the whole genome. We also show that the highly motile cells with IS5 upstream of *flhD*⁺ have greater biofilm formation, although there is a growth cost due to the energetic burden of the enhanced motility as these highly motile cells have a lower yield in rich medium and reduced growth rate. Functional flagella are required for IS5 insertion upstream of *flhD*⁺ as there was no IS5 insertion upstream of *flhD*⁺ for *flhD*, *flgK* and *motA* mutants, and the mutation is stable. Additionally, the IS5 mutation occurs during biofilm formation, which creates genetic and phenotypic diversity. Hence, the cells appear to ‘sense’ whether motility is feasible before a sub-population undergoes a mutation to become hypermotile; this sensing appears related to the master transcription regulator, FlhDC.

The ISME Journal (2011) 5, 1517–1525; doi:10.1038/ismej.2011.27; published online 10 March 2011

Subject Category: evolutionary genetics

Keywords: evolution; IS5; *flhDC*; motility; transposition

Introduction

Bacteria with flagella, such as *Escherichia coli*, have a competitive advantage for moving toward favorable conditions and for avoiding detrimental environments (Fenchel, 2002). The flagellum is a complex organelle that also has an important role in adhesion, biofilm formation and virulence (Pratt and Kolter, 1998; Wood *et al.*, 2006). However, the operation and synthesis of flagella requires significant energy; for example, synthesis involves more than 50 genes at a cost of 2% of the biosynthetic energy in *E. coli* (Soutourina and Bertin, 2003). Hence, coordinating growth and flagella synthesis is necessary. As a result, flagella synthesis is a highly ordered cascade, and on top of the hierarchy, is the master regulator FlhDC required for the expression of all other genes of the flagella regulon (Liu and Matsumura, 1994). The operon that encodes FlhDC, *flhDC*, is one of the most highly regulated loci in the genome and the target for regulation by many environmental factors (Soutourina and Bertin,

2003). The *flhDC* promoter contains many transcription factor-binding sites, including those for global regulatory proteins, such as H-NS and the catabolite gene activator protein complex (Bertin *et al.*, 1994; Soutourina *et al.*, 1999).

Activating genes by insertion of motile DNA elements has been shown to provide benefits to the host under stress by facilitating adaptations to severe environments (Hall, 1998; Petersen *et al.*, 2002; Zhang and Saier, 2009a). As an example, the well-studied *bgl* operon encoding the gene products for the fermentation of β -glucoside depends on the transposition of insertion elements IS1 or IS5 to become active (Schnetz and Rak, 1992; Hall, 1998). Also, the glycerol utilization operon (*glpFK*) can be activated by IS5 when it is inserted upstream (Zhang and Saier, 2009a). For insertion of IS5 into both the *bgl* and *glpFK* operons, the environment influences the mutation rate. In addition, in biofilms, some cells increase their mutation rate during stress by increasing competence during chronic infections (Ehrlich *et al.*, 2010). Hence, these examples demonstrate a Lamarckian-type evolution (that is, environment-driven mutation) that differs from that of Darwin, which holds that mutation rates should not depend on the environment (Koonin and Wolf, 2009).

In recent years, insertion elements IS1 and IS5 have been identified in some *E. coli* strains that

Correspondence: TK Wood, Department of Chemical Engineering, Texas A & M University, 220 Jack E. Brown Building, 3122 TAMU, College Station, TX 77843-3122, USA.

E-mail: Thomas.Wood@chemmail.tamu.edu

Received 16 December 2010; revised 8 February 2011; accepted 8 February 2011; published online 10 March 2011

activate the promoter of *flhDC* and increase motility (Barker *et al.*, 2004; Gauger *et al.*, 2007); however, these mutations were rejected as cases of directed mutation (Barker *et al.*, 2004). Here, we explored the regulation of *flhDC* by IS5 hopping under different environmental conditions and found that the environment influences the IS5 insertion upstream of *flhDC*. Hence, we provide the first example of Lamarckian-type evolution for an active locus.

Materials and methods

Bacterial strains, plasmids and growth conditions

The bacterial strains and plasmids used in this study are listed in Table 1. All experiments were conducted in Luria–Bertani (LB) medium (Sambrook *et al.*, 1989) or M9C medium (Rodriguez and Tait, 1983) at 37 °C, except where indicated. Kanamycin (50 µg ml⁻¹) was used for culturing strains with single-gene knockouts from the Keio Collection (Baba *et al.*, 2006), chloramphenicol (30 µg ml⁻¹) was used for maintaining pCA24N-based plasmids, and ampicillin (100 µg ml⁻¹) was used to maintain plasmid pKD46.

Swimming motility assay

Cell motility was examined as described previously on low-salt, soft-agar plates (1% tryptone, 0.25% NaCl and 0.3% agar), where the wild-type BW25113 is motile (González Barrios *et al.*, 2006). High-salt, soft-agar plates were also used, in which wild-type BW25113 is non-motile due to the high salt concentration (Wang *et al.*, 2009).

Curing IS5 from *flhD*⁺ Ω IS5 by a two-step method

Owing to the stable residence of IS5 after insertion into the *flhDC* regulatory region, curing IS5 via natural excision was not successful. Instead, we used a two-step method to replace the *flhDC* regulatory region of *flhD*⁺ Ω IS5 with that of the

wild-type strain. To facilitate the screening for the right replacement, we first deleted *flhD* from *flhD*⁺ Ω IS5 using P1 transduction to transfer the Δ *flhD* Km^R mutation from BW25113 *flhD* (Table 1) to make a non-motile strain, BW25113 *flhD* Ω IS5; this strain was verified by its complete lack of motility on low-salt, soft-agar plates. Then, the IS5 element was removed and the *flhD*⁺ gene was restored using the one-step inactivation method with pKD46 (Datsenko and Wanner, 2000) by replacing the *flhD* Ω IS5 region with PCR products amplified from the BW25113 wild-type genomic DNA covering the IS5 insertion site and the whole coding region of *flhD*⁺ using primer set PflhDC-4 (PflhDC-F4 and PflhDC-R4) (Table 2). Transformed cells were inoculated into low-salt, soft-agar plates, and cells with restored motility were selected and sequenced to verify the restoration of the wild-type *flhD*⁺ gene and its promoter region.

Crystal violet biofilm assay

Biofilm formation was assayed in 96-well polystyrene plates using 0.1% crystal violet staining (Corning Costar, Cambridge, MA, USA) (Fletcher, 1977). Briefly, each well was inoculated at an initial turbidity at 600 nm of 0.05 and grown without shaking for 8 h in LB and M9C medium. Biofilm formation was normalized by the bacterial growth for each strain (turbidity at 620 nm). Two independent cultures were used for each strain.

Glass wool biofilms

Overnight cultures of the wild-type strain were inoculated into 250 ml LB with 10 g glass wool (Corning Glass Works, Corning, NY, USA) (Ren *et al.*, 2004) to reach a final turbidity 0.05 (600 nm). The cultures were incubated for 8 h, 12 h, 15 h, 24 h and 39 h at 37 °C during which time the cells formed a biofilm on the surface of the glass wool.

Table 1 *Escherichia coli* bacterial strains and plasmids used in this study

Strains and plasmids	Genotype/relevant characteristics	Source
BW25113 strains		
wild type	<i>lac</i> ^R <i>rrnB</i> _{T14} Δ <i>lacZ</i> _{WJ16} <i>hsdR514</i> Δ <i>araBAD</i> _{AH33} <i>ArhaBAD</i> _{LD78}	Baba <i>et al.</i> , 2006
<i>flhD</i> ⁺ Ω IS5	IS5 inserted upstream of <i>flhDC</i> in the opposite orientation in BW25113	This study
<i>flhD</i>	Δ <i>flhD</i> Ω Km ^R , replacement of <i>flhD</i> coding region with Km ^R	Baba <i>et al.</i> , 2006
<i>flhD</i> Ω IS5	IS5 inserted upstream of Δ <i>flhD</i> in the opposite orientation in Δ <i>flhD</i> Ω Km ^R	This study
<i>flhD</i> ⁺ Δ IS5	IS5 removed from <i>flhD</i> ⁺ Ω IS5	This study
<i>flgK</i>	Δ <i>flgK</i> Ω Km ^R , replacement of the <i>flgK</i> coding region with Km ^R	Baba <i>et al.</i> , 2006
<i>motA</i>	Δ <i>motA</i> Ω Km ^R , replacement of the <i>motA</i> coding region with Km ^R	Baba <i>et al.</i> , 2006
<i>hns</i>	Δ <i>hns</i> Ω Km ^R , replacement of the <i>hns</i> coding region with Km ^R	Baba <i>et al.</i> , 2006
<i>hns flhD</i> ⁺ Ω IS5	IS5 inserted upstream of <i>flhDC</i> in the opposite orientation in Δ <i>hns</i> Ω Km ^R	This study
Plasmids		
pCA24N	Cm ^R ; <i>lac</i> ^R	Kitagawa <i>et al.</i> , 2005
pCA24N- <i>hns</i>	Cm ^R ; <i>lac</i> ^R , P _{T5-lac} :: <i>hns</i> ⁺	Kitagawa <i>et al.</i> , 2005
pKD46	Amp ^R , λ Red recombinase expression	Datsenko and Wanner, 2000

Km^R, Cm^R, and Amp^R are kanamycin, chloramphenicol and ampicillin resistance, respectively.

Table 2 Oligonucleotides used for PCR and qPCR in this study

Primer set name	Primer name	Sequence (5'-3')	Source
PflhDC	PflhDC-F2	CCTGTTTCATTTTTGCTTGCTAGC	Barker <i>et al.</i> , 2004
	PflhDC-R3	GGCTGCGATTTTCAATAATGC	This study
PflhDC Ω IS5	PflhDC-F2	CCTGTTTCATTTTTGCTTGCTAGC	Barker <i>et al.</i> , 2004
	PflhDC-IS5-R	CCTGTTCTATGGCTCCAGATGAC	This study
PflhDC-4	PflhDC-F4	TCGACCTCACTTGAGGCAATAA	This study
	PflhDC-R4	TTTCGTGCGTCTCAATCTCTTC	This study
purA	purA-F	GGGCTGCTTATGAAGATAAAGT	Wang <i>et al.</i> , 2009
	purA-R	TCAACCACCATAGAAGTCAGGAT	Wang <i>et al.</i> , 2009
metG	metG-F	GGTGAAGCGCTCTAAAGAAGAAG	This study
	metG-R	AGCAGTTTGTGAGAACCTTCAAC	This study
IS5-1	IS5-F1	CGTGAATAACATCGCCAGTTGGTT	This study
	IS5-R1	CCGGCAAGGTAAGAACCTTGAAAC	This study
IS5-2	IS5-F1	CGTGAATAACATCGCCAGTTGGTT	This study
	IS5-R2	CAAGAACAAAACGGCCATCAACAT	This study
Pade	Pade-F	ATTTGCGGGTTCACAAAAAC	This study
	Pade-R	GGGAAACGGCTAACAATTC	This study
Pbgl	PbglGFB-F	GCGATGAGCTGGATAAACTGCTG	Zhang and Saier, 2009a
	PbglGFB-R	ACTTGAGTTAATCTTTTCGCCAGC	Zhang and Saier, 2009a
Pfuc	PfucAO-F	GCTCTGTAATATGACGGCGGTC	Zhang and Saier, 2009a
	PfucAO-R	AGGCAAGTGTCATAATCTGACGAGC	Zhang and Saier, 2009a
Pglp	PglpFK-F	AAATCCTCGTCCCGATTACC	This study
	PglpFK-R	GGACCTCCACGATGCTTGTA	This study

F indicates the forward primer and R indicates the reverse primer.

At each time point, planktonic cells in contact with biofilms were taken directly from the culture. Glass wool was taken from the culture and was rinsed twice with 100 ml 0.85% NaCl, and biofilm cells were collected after 2 min of sonication of the glass wool with 200 ml 0.85% NaCl in a bath sonicator (Fisher Scientific, Pittsburgh, PA, USA). A population of $\sim 10^8$ cells (OD ~ 1) was inoculated onto a soft-agar plate for the motility assay.

Quantitative PCR (qPCR)

qPCR was performed using the StepOne Real-Time PCR System (Applied Biosystems, Foster City, CA, USA). In all, 100–200 ng of genomic DNA was used for the qPCR reaction using the Power SYBR Green PCR Master Mix (Applied Biosystems). Primers were designed using Primer3Input Software (v. 0.4.0; Rozen and Skaletsky, 2000) and are listed in Table 2. The number of chromosomes that lack IS5 upstream of *flhD*⁺ was quantified using primer set PflhDC (PflhDC-F2 and PflhDC-R3), and the number of chromosomes with IS5 upstream of *flhD*⁺ was quantified using primer set PflhDC Ω IS5 (PflhDC-F2 and PflhDC-IS5-R). To quantify the total copy number of IS5 elements in the chromosome under different growth conditions, the chromosome number was quantified using primer sets purA and metG (Table 2), which amplify single-copy genes *purA* and *metG*, respectively, and the copy number of IS5 was quantified using primer sets IS5-1 and IS5-2 (Table 2), which amplify two different regions of the IS5 gene. The binding efficiencies of the six sets of primers were tested by varying concentrations of templates to generate a standard curve (Pfaffl, 2001), and the templates used were genomic

DNA of the wild-type strain for primer sets PflhDC, purA, metG, IS5-1 and IS5-2, and genomic DNA of the *flhD*⁺ Ω IS5 strain for primer set PflhDC Ω IS5.

Results

IS5 hops upstream of flhD⁺ only when cells swim on motility plates

While performing motility assays using a wild-type *E. coli* K12 BW25113 strain (which has low motility), we noticed that a temporal change occurs in the swimming pattern. When the wild-type cells were plated onto soft-agar plates (0.3% agar) after 8–12 h of incubation, a halo of 11–14 mm in diameter formed (Figure 1, Halo 1). Prolonged incubation (after 24 h) led to some cells swimming faster and forming a second halo with less cell density (Figure 1, Halo 2). The two halos were separated by a zone with very low cell density. Cells from Halo 2 were collected and sequenced in the regulatory region of *flhDC*; the sequencing results (GenBank accession JF342359) showed that an IS5 insertion element (1195 bp, accession J01735) (Kröger and Hobom, 1982) was upstream of *flhD*⁺ in these cells, but not for cells inside the original halo. IS5 was inserted in the opposite orientation into a 4-bp target site (5'-TTAA-3') 96–99 bp upstream of the transcription start site of the *flhDC* operon, the same position described previously for the MG1655 *fnr* strain (Barker *et al.*, 2004). These cells were named *flhD*⁺ Ω IS5. No insertion element was present upstream of *flhD*⁺ for the culture used to inoculate the soft-agar plate. Hence, on soft-agar plates, IS5 hops into the promoter region of *flhDC*.




Figure 1 Swimming halo formed after 12 h (left) and 24 h (right) at 37 °C on low-salt, soft-agar plates. Halo 1 refers to the original halo formed by wild-type swimming cells, and Halo 2 refers to the outside halo formed by *flhD*⁺ Ω IS5 swimming cells.

To quantify the proportion of cells that have acquired IS5 during migration on motility plates, qPCR was conducted using a forward primer that binds upstream of the IS5 insertion site and to one of two reverse primers that bind either to the downstream chromosome region without IS5 or bind inside IS5. The copy numbers obtained from the two separate qPCR reactions were used to estimate the percentage of cells that acquired IS5 upstream of *flhD*⁺. Genomic DNA was isolated using cells collected from different regions of the swimming zone, and was used as templates for qPCR. Starting with a BW25113 wild-type colony (lacks IS5 upstream of *flhD*⁺), in the first stage of migration (less than 12 h) on soft-agar plates (Figure 1, Halo 1), the proportion of cells with IS5 upstream of *flhD*⁺ was very low ($\sim 10^{-7}$), which is similar to that of the culture used for inoculation. During the development of the second halo, $9 \pm 4\%$ cells acquired IS5 after 24 h, and $21 \pm 1\%$ cells acquired IS5 after 36 h. Therefore, within 24 h, a substantial fraction of cells within the second halo had IS5 upstream of *flhD*⁺.

The frequency of IS5 hopping was also tested on plates with the same nutrient content as soft-agar plates, but with higher amounts of agar (1.5%), which serves to immobilize the *E. coli* cells, and on regular LB-agar plates with more nutrients (1.5% agar), which also does not support motility (Copeland and Weibel, 2009). In both cases, after 12 h, 24 h, 48 h and 72 h, the frequency of IS5 hopping upstream of *flhD*⁺ was unchanged and remained very low ($\sim 10^{-7}$, estimated by qPCR). Thus, there was a 10^4 -fold induction of IS5 hopping upstream of *flhD*⁺ on soft-agar plates compared with IS5 hopping on hard-agar plates. In contrast, when cultured in LB liquid medium or in minimal medium, the frequency of IS5 hopping upstream of *flhD*⁺ for planktonic cells remained low ($\sim 10^{-7}$) and did not change for 11 days (~ 200 generations in LB and ~ 100 generations in minimal medium). This occurrence of a motile sub-population of cells was also reported for MG1655 *fnr* from an inoculum of poorly motile MG1655 *fnr* on soft-agar plates, and PCR screening confirmed that IS5 inserted into the *flhDC* regulatory region in the same manner (Barker

et al., 2004). The sequenced K12 strain, MG1655, is a relative of BW25113 (Baba *et al.*, 2006), and is highly motile because of a similar IS1 insertion sequence upstream of *flhD*⁺ (Blattner *et al.*, 1997; Gauger *et al.*, 2007).

Clearly, IS5 hopping upstream of *flhD*⁺ occurred at very different frequencies on plates with different agar concentrations. It is well known that for soft-agar plates (agar <0.4%) the pore size of the gel is larger than the bacterial cells, allowing them to penetrate into the polymer network and swim (Copeland and Weibel, 2009). While on hard-agar plates (agar >1.5%), cells grow into non-motile colonies and retain their planktonic morphology and phenotype. Cells isolated from the edge of the soft-agar plate have distinctively more flagella than cells from a hard-agar plate (Copeland and Weibel, 2009). Increased motility is thus an advantage on soft-agar plates because it allows cells to reach more nutrients. Hence, we conclude that soft-agar plates serve as a type of environmental cue, which promote cell motility by inducing IS5 hopping into the upstream region of *flhD*⁺.

IS5 does not hop upstream of flhD⁺ in flhD and flagella structure mutants

To determine whether IS5 hopping is dependent on the ability to swim, we plated cells with an *flhD* deletion, but which contain the upstream regulatory region and insertion site for IS5. As expected, the *flhD* cells did not swim or form a halo on motility plates after 24 h, 48 h or 72 h incubation. All cells were collected from soft-agar plates and checked for the presence of IS5 by qPCR, and no cells had IS5 in the regulatory region of *flhDC*. Hence, motility is required for IS5 insertion into the regulatory region of *flhDC*.

To determine whether IS5 insertion relies on FlhD or relies on an active flagella component and rotor, the frequency of IS hopping was also tested for strains that lack essential components of the flagella apparatus. MotA is one of the two membrane proteins that form the stator of the flagellar motor (Dean *et al.*, 1984), and FlgK is a hook-associated protein (Komeda *et al.*, 1978). Critically, after 24 h and 36 h of incubation on low-salt, soft-agar plates, no IS5 was detected in strains with these two mutations. Hence, we conclude that IS5 hopping into the regulatory region of *flhDC* requires active motility.

This conclusion is corroborated by results with mutations in the global regulator H-NS (Soutourina *et al.*, 1999). When we plated the *hns* mutant on soft-agar plates, as expected, the *hns* mutant was non-motile due to a reduced expression of *flhDC* and a subsequent lack of flagella biosynthesis (Bertin *et al.*, 1994; Soutourina *et al.*, 1999). However, after long incubation times (24 h) with the *hns* mutant, different from the non-motile *flhD*, *flgK* and *motA* mutants, a halo was formed that contained highly




Figure 2 (a) Early biofilm formation (8 h) at 37 °C in LB medium and M9C medium for the wild-type strain (white bar), strain *flhD*⁺ Ω IS5 that acquired IS5 upstream of the *flhDC* operon (gray bar), and strain *flhD*⁺ Δ IS5 in which IS5 was cured from strain *flhD*⁺ Ω IS5 (slash marks). Data are the average of 10 replicate wells from two independent cultures, and one standard deviation is shown. (b) Swimming motility of the three strains on low-salt, soft-agar plates after 12 h incubation at 37 °C. Three independent cultures were tested, and one representative image is shown. (c) Growth of the wild-type strain (open white circle), *flhD*⁺ Ω IS5 (filled gray circle) and *flhD*⁺ Δ IS5 (filled black triangle) in LB medium at 37 °C. Two independent cultures were used.

motile cells. These cells were found to have the IS5 insertion upstream of *flhD*⁺. Hence, non-structural mutations in the motility regulon permit IS5 hopping into the upstream region of *flhD*⁺ on agar plates that promote motility.

IS5 insertion upstream of flhD⁺ increases motility and biofilm formation while it decreases growth

The *flhD*⁺ Ω IS5 cells formed significantly more biofilm after 8 h than wild-type cells (5 ± 1 -fold more, Figure 2a) and have much greater motility (7 ± 2 -fold greater more, Figure 2b). All cells from four different spots of the outside halo (Figure 1, Halo 2) had the same IS5 insertion upstream of *flhD*⁺ and had the same motility. The specific growth rate of the cells with IS5 in *flhD* was slightly lower, and the cells had a lower yield ($6 \pm 1\%$ reduction) (Figure 2c). This result is consistent with an earlier study, which shows that MG1655 (IS1 inserted upstream of *flhDC*) grew more slowly than MG1655 that lacks IS1 upstream of *flhDC* due to the reduction in energy available for growth due to the enhanced motility (Gauger *et al.*, 2007). For our work, when IS5 was cured from the upstream region of *flhD*⁺, growth, motility and biofilm formation were restored to the wild-type values (Figure 2), which demonstrates that the increase in motility, the increase in early biofilm formation, and the decrease in growth are due to the presence of IS5 in promoter of *flhDC*.

IS5 insertion upstream of flhD⁺ is stable and heritable
The stability of IS5 upstream of *flhD*⁺ was tested in planktonic cells harboring IS5 upstream of *flhD*⁺ using serial dilutions every 24 h for 11 days in LB medium and in minimal medium (M9C). On the basis of qPCR, IS5 is not removed from the upstream of *flhD*⁺ after 1, 5 and 11 days. This result indicates that once IS5 inserted into *flhDC*, it is stable and is not removed (it may make copies and insert into other areas of the chromosome, but it leaves a copy of itself). Moreover, wild-type cells were also assayed using the same conditions, and no cells acquired IS5 in the *flhDC* regulatory region after 1, 5 and 11 days. These results provide another line of evidence that IS5 hopping upstream of *flhD*⁺ occurs only in a motility-inducing environment.

IS5 insertion upstream of flhD⁺ increases diversity in biofilm cells

Flagella are important for initial attachment in biofilms (Pratt and Kolter, 1998), for the mature structure (Wood *et al.*, 2006), and for the dispersal of biofilms (Kaplan, 2010). Here, we tested whether IS5 is inserted upstream of *flhD*⁺ during biofilm development by screening biofilm cells for increases in motility. In the presence of glass wool, planktonic cells were collected from the culture, and biofilm cells were collected from the glass wool by sonicating. Motility results indicated that a small proportion of biofilm cells (as early as 8 h) migrated much faster than the rest of the non-motile cells, which




Figure 3 Emergence of highly motile cells (*flhD*⁺ Ω IS5) during biofilm development. (a) Biofilm (left) and planktonic cells in contact with biofilms (right) were collected at different time points from cultures with glass wool and plated on high-salt, soft-agar plates and incubated for 12 h. Cells with high motility were identified at these stages by the formation of swimming halos. (b) Screening of biofilm cells with high motility after 24 h with high-salt, soft-agar plates where cells were non-motile, unless IS5 was inserted upstream of *flhD*⁺. A total of 500 cells were tested, and only 108 cells are shown. (c) Emergence of highly motile cells (*flhD*⁺ Ω IS5) in biofilms at different stages in 96-well polystyrene plates. Planktonic cells in contact with biofilms (left) and biofilm cells (right) were collected at different time points and plated on high-salt, soft-agar plates and incubated for 12 h. Cells with high motility were identified at these stages by the formation of swimming halos.

appeared as a bright dot in the inoculum ring (Figure 3a, left side of the plate). For the planktonic cells in contact with biofilms, a small proportion of highly-motile cells emerged after 15 h (Figure 3a); however, these motile cells are probably derived from the biofilms as no highly motile cells arise in shaking flasks without biofilms, and cells frequently detach from biofilms (Kaplan, 2010). After 15 h, 24 h and 39 h of incubation, the proportion of highly motile cells increased for both biofilm and planktonic cells in contact with biofilms. A total of 500 cells from biofilms after 24 h were screened on high-salt, soft-agar plates, and 2.2% of the cells had increased motility (Figure 3b). PCR screening for the insertion element upstream of *flhDC* for these cells showed that IS5 was inserted upstream of *flhDC*. Similar results were obtained for cells grown statically in 96-well polystyrene plates; cells with higher motility were found after 12 h of incubation (Figure 3c). Thus, we conclude that IS5 hopping in *flhDC* occurs during biofilm formation and creates a sub-population of cells with increased motility but slower growth.

IS5 hops upstream of flhD⁺, but not into other operons on motility plates and in biofilms

IS5 is present in multiple (10–23) copies in the chromosome of *E. coli* K12 (Deonier, 1996), and in

K12 strain W3110, which is closely related to BW25113 (Jensen, 1993), IS5 is present at 23 copies (Umeda and Ohtsubo, 1990). To test whether the transposition of IS5 upstream of *flhD*⁺ is specific or due to a general increase in IS5 transposition during swimming, the copy number of IS5 in the chromosome was quantified by qPCR and compared with the copy number of *purA* and *metG*, which are present as single copies. For *flhD*⁺ Ω IS5 cells collected from outside halos on soft-agar plates, the estimated copy number of IS5 was 23.6 ± 0.7 , and for BW25113 wild-type cells growing planktonically, there were 22.5 ± 0.8 copies of IS5. Therefore, the highly motile cells have only one additional IS5 element.

Moreover, upstream regions of *bgl*, *ade*, *fuc* and *glpFK* operons were also screened for the presence of insertion elements for *flhD*⁺ Ω IS5 cells collected from outside halos and for *flhD*⁺ Ω IS5 cells collected from 24 h biofilms using PCR screening around the insertion sites (four pairs of primers list in Table 2). Insertion of IS5 or IS1 elements into these operons that derepress gene expression have been reported under specific conditions (Hall, 1998; Petersen *et al.*, 2002; Zhang *et al.*, 2010). As expected, based on the qPCR results, no IS element was found in any of these operons that corroborates that IS5 insertions occurred only upstream of *flhD*⁺. In addition, we also quantified the copy number of

IS5 for all BW25113 cells collected from soft-agar plates after 12 h or 24 h incubation to obtain a population average, and the increase in copy number of IS5 in the population ranged from 0.3 to 1.5 when compared with cells growing planktonically after 24 h in LB, which corroborates again that the IS5 insertions occurred only upstream of *flhD*⁺.

IS5 hopping disrupts H-NS binding to the flhDC operon
In vitro transcription experiments showed that H-NS represses *flhDC* transcription at the AT-rich region that contains the insertion site for IS5 (5'-TTAA-3') in cells without IS5 upstream of *flhDC* (Soutourina *et al.*, 1999). We hypothesized that once IS5 is inserted into this region, it may lead to a loss of repression of *flhDC* by H-NS due to the disruption of the AT-rich region. As expected, we found that H-NS significantly repressed motility when overproduced in the wild-type strain without IS5 upstream of *flhD*⁺, which agrees with earlier studies (Soutourina *et al.*, 1999). However, this repression of motility by H-NS was abolished in cells that had IS5 upstream of *flhD*⁺, for example, in *flhD*⁺ Ω IS5 and *hns flhD*⁺ Ω IS5. Hence, after its insertion, IS5 prevents H-NS from repressing motility.

Discussion

Here, we show that IS5 hopping into the promoter region of *flhDC* occurs at a high frequency only under environmental conditions that promote high flagella activity. It is well known that most transposition is not random, and hot spots have been identified where transposition events occur more frequently. IS5 has been found to be present in multiple (10–23) copies in the chromosome of *E. coli* K12 (Deonier, 1996) and shows a preference for 5'-YTAR-3' (Y = C or T) (R = A or G) target sequences (Mahillon and Chandler, 1998). Upstream of the *flhDC* operon, there is one hot spot (5'-TTAA-3') for IS5 insertion (Barker *et al.*, 2004). However, we provide evidence that this IS5 hopping event is not solely due to the presence of a hot spot for IS5 in the upstream of *flhDC* because prolonged incubation of a mutated strain without *flhD*, *flgK* or *motA* did not increase IS5 hopping even though the hot spot upstream of *flhDC* remains for these three mutants. Another line of evidence is that prolonged incubation leads to increased IS5 hopping on soft-agar plates (swimming is allowed), but not on hard-agar plates (swimming is inhibited).

Once inoculated on agar plates, cells replicate, causing an increase in the population and cell density, until nutrients and resources become limited. For cells on soft-agar plates, when the cell density becomes limited by nutrients and wastes, the highly motile cells have an advantage when they migrate further to obtain nutrients and areas with reduced waste concentrations. Hence, for our

experiments, a starvation condition led to IS5 hopping, which allowed cells to obtain a better environment. Thus, we reasoned that the increased transposition occurs when increased motility leads to an increase in fitness, and we propose that transposition of IS5 upstream of *flhD*⁺ represents a quasi-Lamarckian phenomenon, because the induced genetic change is beneficial to the organism. Just as Koonin and Wolf (2009) found for the prokaryotic CRISPR-cas system, the lines of evidence for IS5 hopping in a quasi-Lamarckian way rather than a Darwinian way are as follows: (i) hopping occurs in a motility-driven environment (soft agar) that leads to the insertion of the motile element, (ii) the resulting modification directly affects motility, which is the same cue that caused the modification, and (iii) the modification is adaptive and is inherited by the progeny of the cell that encounters the mobile element.

As determined by qPCR, the number of pre-existing, highly motile mutants in liquid culture was very low (less than $\sim 10^{-7}$) in a total of 10^7 – 10^8 wild-type cells that were used to inoculate the soft-agar plates. Clearly, selection of pre-existing mutants alone would not be expected to produce 10^4 more cells with high motility, which supports our conclusion that environment-induced mutations contribute more substantially to bacterial evolution under these conditions compared with selection of random pre-existing mutator strains. Moreover, general IS5 transposition was not induced on the soft-agar plates, as the increase in IS5 copy number was limited to basically one IS5 transposition per cell, under these conditions. However, the increase in the proportion of *flhD*⁺ Ω IS5 cells from 24 to 36 h on the soft-agar plates may be due to selective amplification of these highly motile cells.

IS5 has also been shown to behave as a transcriptional enhancer of the otherwise cryptic *E. coli* *bgl* operon to allow cells to utilize β -glucoside (Schnetz and Rak, 1992). IS5 is also capable of activating a metabolic operon *glpFK* to allow cells to utilize glycerol in the absence of the catabolite gene activator protein complex (Zhang and Saier, 2009b). In addition, the insertion of IS element in the cryptic *ade* promoter region results in relief of the H-NS-mediated silencing of *ade* operon, which allows the cells to utilize adenine as the sole source of purines (Petersen *et al.*, 2002). However, in contrast to IS transposition into the *ade*, *bgl* and *glpFK* operons, expression of *flhDC* was not silenced before the transposition, and cells were motile in the absence of IS5 in its promoter (Figure 1, Halo 1); hence, the directed mutation here is fundamentally different in that the transcription of an active gene was increased.

In biofilms, we show that the transposition of IS5 upstream of *flhD*⁺ helps to activate *flhDC* expression and generates a proportion of cells with higher motility and slower growth. Consistent with an earlier study with *Pseudomonas aeruginosa*, where

biofilm-grown cells exhibited more variation in swimming motility than did those from the inoculum (Boles *et al.*, 2004; Boles and Singh, 2008), we also observed cells with increased motility and reduced growth arising in *E. coli* biofilms. Genetic diversity has been proposed as a beneficial feature of biofilms and has been reported in various species (Boles *et al.*, 2004; Allegrucci and Sauer, 2007). The differences in swimming motility phenotypes found in *P. aeruginosa* biofilms were heritable and not produced by planktonic growth, and were dependent on RecA function (Boles *et al.*, 2004). Here, we show that increased variation in motility and growth inside biofilms is caused by transposition of insertion elements, which results in a heritable change. Therefore, IS hopping is a novel mechanism for the production of genetic and phenotypic variations in biofilms.

We have shown previously that phage elements have an important role in generating diversity inside *E. coli* biofilms (Wang *et al.*, 2009), and excision of e14, CP4-57, *rac* and CPS-53 prophages generate isogenetic strains with different phenotypes, including those of motility and growth (Wang *et al.*, 2010). Induction of transposition in prokaryotes under cell-stress conditions is potentially important in creating diversity facilitating adaptation to stressful environments (Hall, 1998; Petersen *et al.*, 2002; Zhang and Saier, 2009a), and IS5 hopping upstream of *flhD*⁺ is an adaptive mutation. Moreover, as IS5 is also found in *E. coli* bacteriophages (Kröger and Hobom, 1982), horizontal gene transfer might influence dissemination of IS5 elements to different strains of *E. coli*. Hence, horizontal gene transfer appears to be a form of quasi-Lamarckian inheritance (Koonin and Wolf, 2009) in prokaryotes, in which transferred genes confer selective advantages for growth in that environment. Here, we show clearly that cells can become highly motile to adapt to starvation conditions due to IS5 hopping and perhaps this is facilitated by horizontal gene transfer.

Acknowledgements

This work was supported by the NIH (R01 GM089999). We are grateful for the Keio and ASKA strains provided by the Genome Analysis Project in Japan. TW is the T Michael O'Connor Endowed Professor at Texas A & M University.

References

- Allegrucci M, Sauer K. (2007). Characterization of colony morphology variants isolated from *Streptococcus pneumoniae* biofilms. *J Bacteriol* **189**: 2030–2038.
- Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M *et al.* (2006). Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection. *Mol Syst Biol* **2**: 0008.
- Barker CS, Pruss BM, Matsumura P. (2004). Increased motility of *Escherichia coli* by insertion sequence element integration into the regulatory region of the *flhD* operon. *J Bacteriol* **186**: 7529–7537.
- Bertin P, Terao E, Lee EH, Lejeune P, Colson C, Danchin A *et al.* (1994). The H-NS protein is involved in the biogenesis of flagella in *Escherichia coli*. *J Bacteriol* **176**: 5537–5540.
- Blattner FR, Guy Plunkett I, Bloch CA, Perna NT, Burland V, Riley M *et al.* (1997). The complete genome sequence of *Escherichia coli* K-12. *Science* **277**: 1453–1462.
- Boles BR, Singh PK. (2008). Endogenous oxidative stress produces diversity and adaptability in biofilm communities. *Proc Natl Acad Sci USA* **105**: 12503–12508.
- Boles BR, Thoendel M, Singh PK. (2004). Self-generated diversity produces ‘insurance effects’ in biofilm communities. *Proc Natl Acad Sci USA* **101**: 16630–16635.
- Copeland MF, Weibel DB. (2009). Bacterial swarming: a model system for studying dynamic self-assembly. *Soft Matter* **5**: 1174–1187.
- Datsenko KA, Wanner BL. (2000). One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci USA* **97**: 6640–6645.
- Dean GE, Macnab RM, Stader J, Matsumura P, Burks C. (1984). Gene sequence and predicted amino acid sequence of the *motA* protein, a membrane-associated protein required for flagellar rotation in *Escherichia coli*. *J Bacteriol* **159**: 991–999.
- Deonier RC. (1996). Native insertion sequence elements: locations, distributions, and sequence relationships. In: Neidhardt FC, Curtiss RI, Ingraham JL, Lin ECC, Low K, Magasanik BB *et al.* (eds). *Escherichia coli and Salmonella: Cellular and Molecular Biology*. 2nd edn. ASM Press: Washington, DC, pp 2000–2011.
- Ehrlich GD, Ahmed A, Earl J, Hiller NL, Costerton JW, Stoodley P *et al.* (2010). The distributed genome hypothesis as a rubric for understanding evolution *in situ* during chronic bacterial biofilm infectious processes. *FEMS Immunol Med Microbiol* **59**: 269–279.
- Fenchel T. (2002). Microbial behavior in a heterogeneous world. *Science* **296**: 1068–1071.
- Fletcher M. (1977). The effects of culture concentration and age, time, and temperature on bacterial attachment to polystyrene. *Can J Microbiol* **23**: 1–6.
- Gauger EJ, Leatham MP, Mercado-Lubo R, Laux DC, Conway T, Cohen PS. (2007). Role of motility and the *flhDC* operon in *Escherichia coli* MG1655 colonization of the mouse intestine. *Infect Immun* **75**: 3315–3324.
- González Barrios AF, Zuo R, Ren D, Wood TK. (2006). Hha, YbaJ, and OmpA regulate *Escherichia coli* K12 biofilm formation and conjugation plasmids abolish motility. *Biotechnol Bioeng* **93**: 188–200.
- Hall BG. (1998). Activation of the *bgl* operon by adaptive mutation. *Mol Biol Evol* **15**: 1–5.
- Jensen KF. (1993). The *Escherichia coli* K-12 ‘wild types’ W3110 and MG1655 have an *rph* frameshift mutation that leads to pyrimidine starvation due to low *pyrE* expression levels. *J Bacteriol* **175**: 3401–3407.
- Kaplan JB. (2010). Biofilm dispersal: mechanisms, clinical implications, and potential therapeutic uses. *J Dent Res* **89**: 205–218.
- Kitagawa M, Ara T, Arifuzzaman M, Ioka-Nakamichi T, Inamoto E, Toyonaga H *et al.* (2005). Complete set of ORF clones of *Escherichia coli* ASKA library (a complete set of *E. coli* K-12 ORF archive): unique resources for biological research. *DNA Res* **12**: 291–299.

- Komeda Y, Silverman M, Matsumura P, Simon M. (1978). Genes for the hook-basal body proteins of the flagellar apparatus in *Escherichia coli*. *J Bacteriol* **134**: 655–667.
- Koonin E, Wolf Y. (2009). Is evolution Darwinian or/and Lamarckian? *Biol Direct* **4**: 42.
- Kröger M, Hobom G. (1982). Structural analysis of insertion sequence IS5. *Nature* **297**: 159–162.
- Liu X, Matsumura P. (1994). The FlhD/FlhC complex, a transcriptional activator of the *Escherichia coli* flagellar class II operons. *J Bacteriol* **176**: 7345–7351.
- Mahillon J, Chandler M. (1998). Insertion sequences. *Microbiol Mol Biol Rev* **62**: 725–774.
- Petersen C, Møller LB, Valentin-Hansen P. (2002). The cryptic adenine deaminase gene of *Escherichia coli*. *J Bio Chem* **277**: 31373–31380.
- Pfaffl MW. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucl Acids Res* **29**: e45.
- Pratt LA, Kolter R. (1998). Genetic analysis of *Escherichia coli* biofilm formation: roles of flagella, motility, chemotaxis and type I pili. *Mol Microbiol* **30**: 285–293.
- Ren D, Bedzyk LA, Thomas SM, Ye RW, Wood TK. (2004). Gene expression in *Escherichia coli* biofilms. *Appl Microbiol Biotechnol* **64**: 515–524.
- Rodriguez RL, Tait RC. (1983). *Recombinant DNA Techniques: An Introduction*. The Benjamin/Cummings Publishing Company: Menlo Park, CA, USA.
- Rozen S, Skaletsky HJ. (2000). Primer3 on the WWW for general users and for biologist programmers. In: Krawetz S, Misener S (eds). *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press: Totowa, NJ, pp 365–386.
- Sambrook J, Fritsch EF, Maniatis T. (1989). *Molecular Cloning: A Laboratory Manual*. 2nd edn. Cold Spring Harbor Laboratory Press: New York, USA.
- Schnetzer K, Rak B. (1992). IS5: a mobile enhancer of transcription in *Escherichia coli*. *Proc Natl Acad Sci USA* **89**: 1244–1248.
- Soutourina O, Kolb A, Krin E, Laurent-Winter C, Rimsky S, Danchin A *et al.* (1999). Multiple control of flagellum biosynthesis in *Escherichia coli*: role of H-NS protein and the cyclic AMP-catabolite activator protein complex in transcription of the *flhDC* master operon. *J Bacteriol* **181**: 7500–7508.
- Soutourina OA, Bertin PN. (2003). Regulation cascade of flagellar expression in Gram-negative bacteria. *FEMS Microbiol Rev* **27**: 505–523.
- Umeda M, Ohtsubo E. (1990). Mapping of insertion element IS5 in the *Escherichia coli* K-12 chromosome: chromosomal rearrangements mediated by IS5. *J Mol Biol* **213**: 229–237.
- Wang X, Kim Y, Ma Q, Hong SH, Pokusaeva K, Sturino JM *et al.* (2010). Cryptic prophages help bacteria cope with adverse environments. *Nat Commun* **1**: 147.
- Wang X, Kim Y, Wood TK. (2009). Control and benefits of CP4-57 prophage excision in *Escherichia coli* biofilms. *ISME J* **3**: 1164–1179.
- Wood TK, Gonzalez Barrios AF, Herzberg M, Lee J. (2006). Motility influences biofilm architecture in *Escherichia coli*. *Appl Microbiol Biotechnol* **72**: 361–367.
- Zhang Z, Saier Jr MH. (2009a). A mechanism of transposon-mediated directed mutation. *Mol Microbiol* **74**: 29–43.
- Zhang Z, Saier Jr MH. (2009b). A novel mechanism of transposon-mediated gene activation. *PLoS Genet* **5**: e1000689.
- Zhang Z, Yen MR, Saier Jr MH. (2010). Precise excision of IS5 from the intergenic region between the *fucPIK* and the *fucAO* operons and mutational control of *fucPIK* operon expression in *Escherichia coli*. *J Bacteriol* **192**: 2013–2019.